

POPULATION ECOLOGY OF TIMBER TREES OF FOOD AND MEDICINAL VALUES IN OBAN FOREST RESERVE, CROSS RIVER STATE, NIGERIA

Olajide^{1*}, O., Annah², A. A. and Etigale¹, E. B.

ABSTRACT

¹Department of Forestry and Natural Environmental Management, University of Uyo, Uyo, Nigeria e-mail: solaolajide1967@gmail.com. ²Department of Forestry and Wildlife, Cross River State University of Technology (Obubura Campus), Obubura, Cross River State, Nigeria *Correspondent Author:

Paucity of quantitative and qualitative ecological data is one of the major factors militating against sustainable natural forest management in Nigeria. Thus, assessment of population density of adult or mature stands and regeneration or juvenile stands of timber-tree species producing important food and medicinal items in Oban Forest Reserve, Cross River State, Nigeria was carried out using transect and quadrat sampling techniques. Results revealed that 19 tree species were encountered with *Coula edulis* having the highest population density (48 per/ha) of adult individuals, while *Cola argentea* was the only species with no adult individual stand but its regeneration were encountered. The highest regeneration population density of 113 per/ha was recorded for *Pycnanthus angolensis*, while the regeneration of three species, viz; *Alstonia boonei*, *Brachystegia eurycoma* and *Rauvolfia vomitoria*, were not encountered but their adult individuals were present in the forest. All the tree species were in a rare status except *C. edulis*, *P. angolensis* and *Vitex* spp. Moreover, only *P. angolensis* had sustainable natural regeneration potential. It is concluded that a ban should be placed on the exploitation of the trees for timber to forestall their extinction from the forest, and enrichment planting should be carried out in the forest using the raised seedlings of the tree species to enhance biodiversity richness.

Keywords: Population ecology, timber trees, medicinal values, Oban forest reserve.

INTRODUCTION

A great number of timber tree species in the natural forests, particularly rainforest, produce various non-timber products which in most cases are much more valuable than wood. The non-timber products are mostly food, medicinal items and raw materials to small and cottage industries. Forest is often erroneously viewed as a crop of merchantable timber trees rather than an interdependent high diversity ecosystem of potential multiple value (Panayotou and Ashton, 1992; Olajide *et al.*, 2010). Until recently, little or no recognition was given to the non-timber products in the scheme of forest management. These products are often ignored in the reckoning of forestry contributions to the gross domestic products (Lorbach *et al.*, 2000). According to Ford Foundation (1998) and Lorbach *et al.* (2000), non-timber forest products are a particularly important part of multiple-use strategies, because they increase the range of income generating options of forest-dependent communities, while avoiding some of the ecological costs of timber cutting. Evaluating from a long-term standpoint, non-timber products from forest trees are much more valuable than their timber as the former can be harvested for many years without cutting down the trees, in contrast to timber harvesting and their harvesting activity or extraction has imperceptible perturbation on the ecosystem (Ella and Domingo, 2014; Udo, 2016).

The predominant natural forest type in Nigeria is rainforest. The Nigerian rainforest is an integral of the world tropical rainforest which has been acknowledged as the most biologically diverse ecosystem on earth (Turner, 2001; Gillespie *et al.*, 2004; Sukhdev, 2010). According to Mgeni (1991), with the unique diversity of plant and animal life, tropical rainforest represents biologically renewable resources of food, medicine, fibre and fuel if well managed. A considerable volume of food items in the forms of fruits seeds, nuts, leaves, twigs, bark, roots and flowers are collected from the rainforest (Ikojo *et al.*, 2003). These food items greatly complement the food supply from cultivated crops. According to Oni and Gbadamosi (1998), the dietary contribution of forest trees to improved nutritional status of mankind is further enhanced by the timing of their availability which often falls at the strategic periods of general food shortage particularly in Nigeria. The tropical rainforest is an outstanding 'natural pharmacy'. According to FAO (1985) and Spore-Dossier (2011), tropical forests provide the only medicine available to the largest proportion of the world population of which 90-95% is in the developing world, and the high concentration of different chemicals in some plants serve as raw materials for the production of modern drugs. The World Health Organization (WHO) has estimated that 80% of the population of sub-sahara Africa relies on plant based medicines for everyday healthcare needs (Bannerman, 1993).

The tropical rainforests have been widely recklessly exploited for timber, and subsequently converted to other forms of land-use aside forestry. Consequently, a considerable genetic resource has been lost, and many species are on the verge of extinction. Among the species threatened by extinction are trees that produce, in addition to timber, food and medicinal products. Unfortunately, most of these trees are not amenable to plantation environment, because of a number of silvicultural and ecological constraints, which include poor viability of seeds, endemic pests and diseases and very slow growth rate (Akinsanmi and Akindele, 2002 and Olajide *et al.*,

2010). Most of these constraints are surmounted while the trees grow in the natural forest ecosystem. It implies that the continuous availability of the food and medicinal products of these trees depends on sustainable management of the remaining areas of the tropical rainforest.

Sustainable management of a tract of natural forest like tropical rainforest would be impossible without quantitative ecological data relating to the population density of adult trees, stem-diameter distribution and regeneration density. This paper is a report of a study on stand density, and regeneration of trees that produce food and medicinal products in a rainforest estate or reserve in Nigeria. It is hoped that the data and information garnered would enhance the conservation of the tree species in particular and sustainable management of the forest estate or reserve in general.

MATERIALS AND METHODS

Study area

The study was carried out in Oban Forest Reserve, Cross River State, Nigeria. The forest is part of the eastern bloc of Nigerian rainforest. The forest reserve covers an area of 245.48 km². The area lies between latitudes 5° 00' and 5° 57' N and longitudes 8° 10' and 8° 55' E. The annual rainfall of the area varies between 2,500mm and 3,500mm, means minimum and maximum annual temperatures are 23 °C and 30°C respectively. The average relative humidity is 85% at 12.00 hour (Offiong and Iwara, 2011). The soil is known to be formed from rocks of pre-cambrian complex especially granite, gneisses and schist (Offiong and Iwara, 2011). The forest reserve has been subjected to timber harvesting or exploitation.

Data collection

One kilometer (1 km) baseline was laid 20 m away from the main access route into the forest. Thereafter, five points of 200 m apart from one and other were marked along the baseline, three points were randomly selected for the laying transects. Thus three 1 km transect were perpendicularly laid to the baseline into the forest. Mature stands (≥ 20 cm diameter at breast height) of individual tree species producing edible and medicinal items within 10 m on both sides of each transect were identified and enumerated by numbering them with indelible red paint to avoid multiple enumerations. Thus, the total sample area assessed for adult or mature trees was 60,000 m² (6 ha). In assessing regeneration or juvenile trees (< 20 cm diameter at breast height), twenty 10 m x 10 m quadrats were randomly alternately laid on both sides of each transect and enumerated for the juvenile individuals of trees of food and medicinal products. Accordingly, a total area of 3,000 m² (0.6 ha) was assessed for juvenile trees. The data collections spanned years 2013 to 2015.

Data analysis

The frequency of adult individual of a tree species was determined in order to estimate its population per hectare by dividing the total population in the sampled area by 6. The density of regeneration or juveniles of each tree species was extrapolated from the population in 0.6 ha, which was the total area enumerated for regeneration or juvenile trees.

RESULTS AND DISCUSSION

The results indicated that 19 timber tree species producing food and medicinal items were encountered (Table 1). The highest population density of adult tree stands (48 per/ha) was recorded for *Coula edulis* while the only species with no adult individual was *Cola argentea* but its regeneration or juvenile stands were encountered (Table 1). The highest regeneration population density of 113 per/ha was recorded for *Pycnanthus angolensis*, while the regeneration of three species, namely, *Alstonia boonei*, *Brachystegia eurycoma* and *Raufofia vomitoria* were not encountered, but they had adult stands (Table 1). Accordingly, the results of adult trees' population density indicated that all the tree species were in rare state except *Coula edulis*, *Pycnanthus angolensis* and *Vitex* spp. (Table 1). The parts of the trees that are extracted for food and medicine differ from species to species, and these include fruits, seeds, nuts, barks, leaves and twigs (Table 2).

According to Parthasarathy and Karthikeyan (1997) and Olajide *et al.* (2015) a tree species with less than 10 individuals per hectare is considered as a rare species. Accordingly, the results on adult trees' population density indicated that all the tree species were in rare state except *Coula edulis*, *Pycnanthus angolensis* and *Vitex* spp (Table 1). Moreover, the results of this study also indicated that only *Pycnanthus angolensis* has sustainable natural regeneration potential. According to Nwoboshi (1982), Dike (1992) and Akpan-Ebe (2005), any tree species in the tropical rainforest with less than one hundred (100) regeneration population per/ha is deemed to have unsustainable natural regeneration potential, because it had been observed that an average of 10 out of 100 regeneration survived and grew up to mature trees. The low population density of most of the tree species reported in this study is similar to the findings of previous workers on population density and regeneration potentials of timber tree species producing economically valuable non-timber products in some areas of rainforest (Ganesan and Davidar, 2003; Olajide *et al.*, 2010 and 2015). The fewer population of individual tree species observed in the study forest may be linked to their over-exploitation for timber. This might have also caused gross inadequacy of seeds for regeneration, as a lot of mother trees (seed producing trees) have been felled for timber, and seeds produced by the few available mother trees are frequently collected as food and medicinal items.

Agyeman (2013) and Newin (2014) reported poor population density of many commercial tree species (including tree producing edible and medicinal fruits and seeds) in a rainforest subjected to timber exploitation in Ghana and Myannar respectively.

Table 1: Population density of timber trees producing food and medicine in Oban Forest Reserve, Cross River State, Nigeria

Tree species	Population density of adult tree (per/ha)	Population density of regeneration (per/ha)
<i>Allanblackia floribundia</i>	1	10
<i>Alstonia boonei</i>	1	-
<i>Baillonella toxisperma</i>	1	8
<i>Brachystegia eurycoma</i>	5	-
<i>Canarium schwinfurthii</i>	1	5
<i>Cola argentea</i>	-	5
<i>Coula edulis</i>	48*	65
<i>Dacryodes edulis</i>	1	5
<i>Dialium guineense</i>	1	22
<i>Enantia chlorantha</i>	5	48
<i>Gambeya albida</i> (formerly <i>Chrysophyllum albidum</i>)	1	30
<i>Irvingia gabonensis</i>	9	32
<i>Pentaclethra macrophylla</i>	2	43
<i>Poga oleosa</i>	1	20
<i>Pycnanthus angolensis</i>	20*	113**
<i>Rauvolfia vomitoria</i>	1	-
<i>Treculia africana</i>	1	23
<i>Vitex spp.</i>	12*	60
<i>Xylopi aethiopica</i>	4	35

*Not-Rare; **Capable of sustainable natural regeneration

Table 2: Part of timber trees used as food and medicine in Oban Forest Reserve, Cross River State, Nigeria

Trees species	Tree part used	Uses
<i>Allanblackia floribundia</i>	Seeds	Food (oil)
<i>Alstonia boonei</i>	Leaves, bark	Medicine
<i>Baillonella toxisperma</i>	Seeds	Stomach ailment; rheumatics pains
<i>Brachystegia eurycoma</i>	Seeds	Food (soup condiment)
<i>Canarium schwinfurthii</i>	Fruit	Food (snacks)
<i>Cola argentea</i>	Fruit mesocarp	Snacks
<i>Coula edulis</i>	Seed (nuts)	Snacks
<i>Dacryodes edulis</i>	Fruit mesocarp	Food (fat/oil)
<i>Dialium guineense</i>	Fruit mesocarp	Snacks
<i>Enantia chlorantha</i>	Bark	Medicine (malaria; jaundice)
<i>Gambeya albida</i> (formerly <i>Chrysophyllum albidum</i>)	Fruit mesocarp	Beverage
<i>Irvingia gabonensis</i>	Fruit mesocarp; seeds	Beverage; soup condiment
<i>Pentaclethra macrophylla</i>	Seeds	Snacks (delicacy)
<i>Poga oleosa</i>	Seeds	Food (oil)
<i>Pycnanthus angolensis</i>	Leaves	Medicine (tooth ache)
<i>Rauvolfia vomitoria</i>	Leaves	Medicine (sedative and tranquilizer; ringworm)
<i>Treculia africana</i>	Seeds	Food (snacks)
<i>Vitex spp.</i>	Fruit	Food (snacks)
<i>Xylopi aethiopica</i>	Fruit & seeds	Spice; medicine (Rheumatic and boil pains)

CONCLUSION AND RECOMMENDATION

The very low population density and unsustainable natural regeneration potentials of most of the tree species require a ban on the exploitation of the trees for timber to forestall their extinction from the forest reserve. Also, enrichment planting operation should be carried out by planting the raised seedlings of the tree species in the forest reserve to boost their stocking, and largely enhance biodiversity richness of the forest.

REFERENCES

- Agyeman, V. K. 2013. The impact of logging damage on tropical rainforest, their recovery and regeneration. *Tropical Forest Update*, 22 (2): 23-25.
- Akinsanmi, F. A. and Akindele, S. O. 2002. Timber yield assessment in the natural forest area of Oluwa forest Reserve, Nigeria. *Nigerian Journal of Forestry*, 32 (1 & 2): 16-22.

- Akpan-Ebe, I. N. 2005. Gap-phase dynamics and natural regeneration in the Stubbs Creek Forest Reserve, Akwa Ibom State, Nigeria. Unpublished PhD Thesis, University of Ibadan, 221p.
- Bannerman, R. H. 1993. Traditional medicine healthcare coverage. W.H.O. Geneva.
- Dike, M. C. 1992. Tree regeneration, recruitment and mortality in Nigeria tropical moist forest. Unpublished Ph.D Thesis, University of Ibadan. 255p.
- Ella, A. B. and Domingo, E. P. 2014. Making the most of NTFPs (Non-Timber Forest Products). *Tropical Forest Update*, 23 (2): 16-17.
- FAO. 1995. Forest, trees and people. Report No. 2, FAO, Rome.
- Ford Foundation 1998. *Forestry for sustainable rural development*. A review of Ford Foundation supported community forestry program in Asia. p.58.
- Ganesan, R. and Davidar, P. 2003. Effect of logging on the structural and regeneration of important fruit bearing trees in a wet evergreen forest, Southern Western Ghats, India. *Journal of Tropical Forest Science*, 15 (1): 12-25.
- Gillespie, T. W., Brock, J. and Wright, C. W. 2004. Prospect of quantifying structure, floristic composition and species richness of tropical forest. *International Journal of Remote Sensing*, 25(14): 70 – 77.
- Ikojo, H. A., Olajide, O. and Asomugha, V. U. 2003. Growth of *Pentaclethra macrophylla* (Benth) seedlings grown on different potting mixture. *Journal of Sustainable Tropical Agricultural Research*, 5: 66-69.
- Lorbach, J., Russo, L. and Vantomme, P. 2000. Needs and constraints for improved inventory and harvesting techniques for non-wood forest products. Seminar proceedings organised by FAO/ECE/ILO. pp. 191-197.
- Mgeni, A. S. M. 1991. A more efficient forest estate: Natural forest or plantations. In Pyatt, N. and Williams, I. (eds.) *Tropical Forestry: Third World priorities versus Western concerns*. University of Wales, Bangor, UK. pp. 43-57.
- Newin, R. 2014. Population dynamics of commercial tree species after selective logging in Myannar. *Tropical Forest Update*, 22(4): 22-25.
- Nwoboshi, L. C. (1982). *Tropical silviculture: Principles and techniques*. Ibadan University Press, Ibadan, Nigeria.
- Offiong, R. A. and Iwara, A. I. 2011. Effects of fallow-genealogical cycles on the build-up of nutrients in soils of Cross River rainforest South-South Nigeria. *Ethiopian Journal of Environmental Studies and Management*, 4: 56-74.
- Olajide, O., Udo, E. S. and Otu, D. O. 2010. Stand density and natural regeneration status of timber tree species producing valuable non-timber products in two tropical rainforests in Southeastern Nigeria. *Journal of Agriculture, Biotechnology and Ecology*, 3 (3): 98-104.
- Olajide, O., Udo, E. S. and Otu, D. O. 2015. Stock density of 'timber plus trees' and their economically valuable non-timber products in Cross River North Forest Reserve, Nigeria. *Journal of Forestry, Environment and Sustainable Development*, 1 (1): 47-51.
- Oni, O. and Gbadamosi, A. F. 1998. Progeny variation in seedlings of *Dacryodes edulis* G. Don. *Journal of Tropical Forest Resources*, 14 (1): 38-47.
- Panayotou, T. and Ashton, P. S. 1992. *Not by timber alone: Economic and ecology for sustaining tropical forests*. Island Press, Washington D.C. and Covelo, California, U.S.A. p. 89.
- Parthasarathy, N. and Karthikeyan, R. 1997. Biodiversity and population density of woody species in a tropical evergreen forest in Courtallum reserve forest, Western Ghats, India. *Tropical Ecology*, 38: 297-306.
- Spore-Dossier 2011. Forest: Conserving the green gold. *Spore*, No. 153: 13-17.
- Sukhdev, P. 2010. The economics of biodiversity and ecosystem services of tropical forests. *Tropical Forest Update*, 20 (1): 8-10.
- Turner, I. M. 2001. *The Ecology of Trees in Tropical Rainforest*. Cambridge University Press, Cambridge. 298p.
- Udo, E. S. 2006. Our forests: Uses, management and abuses. The 51st Inaugural Lecture in the University of Uyo, Uyo, Nigeria. 95p.